

The cracked tooth conundrum: Terminology, classification, diagnosis, and management

WILLIAM KAHLER, MSCDENT, DCLINDENT, FRACDS

ABSTRACT: Purpose: To provide an overview of the clinical features, diagnosis, classification and management of cracked teeth which may be a diagnostic challenge in clinical practice. **Results:** Cracks may initiate from coronal tooth structure or from within the root and affect healthy or root treated teeth. There are many terminologies and classifications in the literature for cracked teeth that can be as confusing as the array of clinical symptoms which are associated with this condition. The term “cracked tooth syndrome” is misleading as there are a range of symptoms that do not form a distinct and reliable pattern. Symptoms will vary with teeth that have healthy pulps, for teeth with inflamed or necrotic pulps, and for teeth that have been root treated. The American Association of Endodontists have classified five specific variations of cracked teeth; craze line, fractured cusp, cracked tooth, split tooth, and vertical root fracture. The importance of differentiating dentin, pulpal and periodontal pain for diagnosis and treatment for these specific entities will be elaborated. A decision flow chart indicating the treatment options available is presented. (*Am J Dent* 2008;21:275-282).

CLINICAL SIGNIFICANCE: A cracked tooth should be considered in the diagnosis of teeth which are sensitive to bite and thermal change. The American Association of Endodontists classification of cracked teeth is useful, though non-vital and root filled cracked teeth and teeth with periapical pathosis should be also considered in forming a diagnosis.

✉: Dr. William Kahler, University of Queensland, Dental School, 200 Turbot St., Brisbane, 4000 Australia. E-✉: w.kahler@uq.edu.au

Introduction

Cracked or incompletely fractured teeth can become symptomatic. Patients often present with a protracted history of pain of varying intensity; the origin of which may be difficult to locate. While intermittent pain on biting is the most consistent complaint associated with these teeth, cracks in teeth may result in a wide range of symptoms ranging from occasional discomfort to severe and prolonged pain. Symptoms are often dependent on the depth and direction of the crack and the tissues involved.

Cracks in teeth may occur in both horizontal and vertical directions involving the crown and/or root. The etiology is generally a result of occlusal forces and iatrogenic procedures.¹ Crown and crown-root fractures are usually incomplete fractures commencing in the crown of posterior teeth from an internal line angle at the floor of a restoration, and often involving a marginal ridge with the fracture extending in a mesiodistal direction. The fracture commences in the crown and may terminate in the vicinity of the cemento-enamel junction or extend apically into the root.²⁻⁶ Vertical root fractures are longitudinally orientated fractures of the root that extend from the root canal to the periodontium.⁷ These fractures are usually complete and extend a variable length along the root generally in a bucco-lingual direction and may extend into the crown.^{4,8-10}

This paper reviewed the literature for an appropriate classification for cracked teeth and to determine the symptoms and processes that allow for correct diagnosis and treatment.

TERMINOLOGY AND DEFINITION

Many authors have proposed different terminologies and definitions for cracks in teeth (Table 1). Gibbs¹¹ first described the clinical symptoms of incomplete fracture of posterior teeth involving the cusp, naming it “cuspal fracture odontalgia”.

Cameron^{2,12} coined the term “cracked tooth syndrome” in describing signs and symptoms associated with cracked teeth. However, there is considerable overlap and confusion in these proposals. For instance, cracked tooth syndrome has been defined as an “incomplete fracture of a vital posterior tooth involving the dentin and possibly the dental pulp”¹³ despite the fact that Cameron¹² reported that only 75% of teeth with “cracked tooth syndrome” will have vital pulps. Many authors confuse the terminology by illustrating teeth with “cracked tooth syndrome” which are in fact teeth with vertical root fractures.¹⁴ The term “cracked tooth syndrome” is misleading as there are a range of symptoms that do not form a distinct and reliable pattern. Symptoms will vary with teeth that have healthy pulps, for teeth with inflamed or necrotic pulps, and for teeth that have been root treated.

Ellis¹⁵ defined incomplete tooth fracture as a “fracture plane of unknown depth and direction passing through tooth structure that, if not already involving, may progress to communicate with the pulp and/or periodontal ligament”. Cracks in teeth can be found in symptomatic and asymptomatic teeth, and are an etiological factor in pulpal disease. This can be a direct result of fracture extension to involve the pulp chamber as bacteria have been reported to be present in cracks,^{16,17} or, indirectly *via* the microleakage of bacterial toxins.¹⁸

CLASSIFICATION

Several authors have proposed classifications which are generally based on either the type or location of the crack, the direction and extent of the crack, and/or the risk of symptoms and/or pathological processes (Table 2).

The American Association of Endodontists, in a document titled “Cracking the Cracked Tooth Code”⁴⁰ identified five types of cracks in teeth which can be viewed at <http://www.aae.org/dentalpro/colleagueneews.htm> and are briefly described in Table 3.

Table 1. Terminology and definitions for cracks/fractures in teeth.

Year	Author(s)	Terminology	Definition
1954	Gibbs ¹¹	Cuspal fracture odontalgia	
1954	Thoma ¹⁹	Fissured fracture	A crack in the crown of the tooth
1957	Ritchey <i>et al</i> ²⁰	Incomplete tooth fracture	
1957	Down ²¹	Fissural fracture	Fractures involving enamel and dentin without loss of tissue
1961	Sutton ²²	Crack lines	A break in the continuity of the tooth revealed only by the presence of a visible transverse line
1962	Sutton ²³	Greenstick fractures	A fracture line forms in a part of a tooth underlying a cusp
1964	Cameron ²	Cracked tooth syndrome	
1972	Wiebusch ²⁴	Hairline fracture	
1973	Hiatt ³	Incomplete crown-root fracture	
1974	Talim & Gohi ²⁵	Incomplete coronal fracture	
1976	Silvestri ²⁶	Split-root syndrome	
1977	Maxwell & Braly ²⁷	Incomplete tooth fracture	A fracture of tooth structure which extends into dentin but in which the tooth remains grossly intact
1981	Andreassen ²⁸	Enamel infraction	An incomplete fracture (crack) of the enamel without loss of tooth substance
1981	Caufield ²⁹	Hairline tooth fracture	
1981	Johnson ³⁰	Crown craze/crack	Injury of enamel without loss of enamel
1983	Abou-Ras ³¹	Tooth structure cracks	A line that breaks or splits the continuity of tooth dentin surface but does not perceptibly separate the surface
1983	Abou-Rass ³¹	Crack/craze lines	Located in coronal enamel
1984	Luebke ³²	Incomplete tooth fracture	A demonstrable fracture but with no visible separation of the segments along the plane of fracture
1984	Kruger ³³	Cracked cusp syndrome	
1986	Brännström ¹⁸	Dentin crack syndrome	
1988	Williams ³⁴	Incomplete vertical tooth fracture	
1989	Lost <i>et al</i> ³⁵	Tooth infraction	
1989	Schweitzer ³⁶	Odontiatrogenic tooth fracture	
1990	Ehrmann & Tyas ¹³	Cracked tooth syndrome	Incomplete fracture of a vital posterior tooth involving the dentin and possibly the dental pulp
1998	Zuckerman ³⁷	Cracked tooth	Fractured segments are still joined to one another by a portion of that tooth through which the fracture has not yet extended
2001	Ellis ¹⁵	Incomplete tooth fracture	A fracture plane of unknown depth and direction passing through tooth structure that, if not already involving, may progress to communicate with the pulp and/or periodontal ligament

Table 3. American Association of Endodontists classification of cracked teeth.

Classification	Originate	Direction	Symptoms	Pulp Status	Prognosis
Craze Line	Crown	Variable	None	Vital	Excellent
Fractured cusp	Crown	M-D and/or F-L	Mild and generally, only to biting and cold	Usually vital	Good
Cracked tooth	Crown±Root	M-D often Central	Acute pain on biting Occasionally sharp pain to cold	Variable	Questionable: Dependent on depth and extent of the crack
Split tooth	Crown+Root	M-D	Marked pain on chewing	Often root filled	Poor unless crack terminates just subgingivally
Vertical root fracture	Roots	F-L	Vague pain Mimics periodontal disease	Mainly root filled	Poor: Root resection in multi-rooted teeth

Craze lines are found in the majority of adult teeth and only involve enamel. In posterior teeth, craze lines are usually evident crossing marginal ridges and/or extending along buccal and lingual surfaces. Long vertical craze lines are often found in anterior teeth (Fig. 1).

Fractured cusps usually result from insufficient cusp support when the marginal ridge is weakened by an intra-coronal restoration (Fig. 2). The crack often extends in mesio-distal and bucco-lingual directions commonly involving one or both marginal ridges as well as a buccal or lingual groove and terminates in the cervical region either parallel to the gingival margin or slightly subgingival.

A cracked tooth is indicative of a crack extending from the occlusal surface of the tooth apically without separation of the two segments. The crack is generally located centrally in a

mesio-distal direction and may involve one or both marginal ridges (Fig. 3).

A split tooth is indicative of a crack extending through both marginal ridges usually in a mesio-distal direction splitting the tooth completely into two separate segments (Figs. 4a-c). The crack is generally located centrally in the tooth and this entity is the result of crack propagation of a cracked tooth.

Vertical root fractures commence in the root generally in a bucco-lingual direction (Figs. 5a-b). The crack is generally complete though may be incomplete and involve only one surface. The crack may involve either the entire root or only a portion of the root.

The American Association of Endodontists classification identifies four types of cracks that are located in the crown-root as well as vertical root fractures that originate from the



Fig. 1. Craze lines in enamel are evident.



Fig. 2. A fractured cusp is located at the base of the cavity (arrow).




Fig. 3. A cracked tooth is shown where there is a mesio-distal crack without separation of the segments.

Table 2. Proposed classifications for cracked teeth.

Pruden³⁸

- A. Crack line
 - 1. No separation of parts, no pain symptoms
 - 2. No apparent separation, but tooth sensitive to percussion-or patient has persistent, vague pain not definitely related to the tooth
- B. Fractured cusp
 - 1. No pain or pulp involvement
 - 2. Possible pulp involvement
- C. Fractured crown
 - 1. No pulpal involvement
 - 2. Pulp involved
- D. Fractured root tip

Talim & Gohil²⁵

- Class 1 - Fracture involving enamel
 - a. Horizontal or oblique
 - b. Vertical
 - 1. Complete
 - 2. Incomplete
- Class 2 - Fracture involving enamel and dentin without involving pulp
 - a. Horizontal or oblique
 - b. Vertical
 - 1. Complete
 - 2. Incomplete
- Class 3 - Fracture of enamel and dentin involving the pulp
 - 1. Horizontal
 - 2. Vertical
 - 1. Complete
 - 2. Incomplete
- Class 4 - Fracture of the roots
 - a. Vertical or oblique
 - 1. Involving the pulp
 - 2. Not involving the pulp
 - b. Horizontal
 - 1. Cervical third
 - 2. Middle third
 - 3. Apical third

Luebke³²

- Class 1 - Incomplete, supra-osseous with no periodontal defect
- Class 2 - Incomplete, intra-osseous with a minor periodontal defect
- Class 3 - Complete or incomplete, intra-osseous with a major periodontal defect

Williams³⁴

- Category 1 - Incomplete vertical fracture through enamel into dentin but not into pulp
- Category 2 - Incomplete crown fracture involving the pulp
- Category 3 - Incomplete vertical fracture crossing the attachment
- Category 4 - Fracture divides the tooth completely

Clark et al³⁹

- Type 1 Cracks - Little or no risk of underlying pathology
- Type 2 Cracks - Moderate risk of underlying pathology
- Type 3 Cracks - High risk of underlying pathology

root. For the purpose of this review, the four coronal fractures will be considered together to assess aspects of tooth fracture such as diagnosis, causes, mechanisms and treatment of fractured teeth.

INCIDENCE

The presence of a cracked tooth occurs primarily in adulthood. Cameron² reported that 80% of 102 cracked teeth occurred with patients over 40 years of age. Other reports^{3,12,31,42-45} about the incidence and prevalence of cracked teeth were commonly associated with intracoronal restorations and most prevalent in mandibular molars. The wedging effect of the prominent mesio-palatal cusp of the maxillary first molar may account for this observation.^{3,43} The transverse ridge of the maxillary molars may provide structural reinforcement and account for the lower incidence of fracture in these teeth.³ The maxillary molars and premolars have a similar incidence of fracture, with the mandibular premolars being the least susceptible.^{30,42-44}

The disto-lingual cusp of mandibular molars is the most susceptible cusp for fracture. The findings for the prevalence of cusp fracture in other teeth were not consistent.^{42,44-45} Non-functional cusps may be more susceptible to fracture than functional cusps.^{42,45} This observation may be a result of cuspal dimension as functional cusps are significantly larger in a bucco-lingual dimension and are covered with a thicker layer of enamel.⁴⁶ While functional cusps are supported on the inner and outer inclines by the opposing tooth, non functional cusps may be more susceptible to fracture from lateral excursive occlusal forces due to the lack of support from the outer incline.⁴⁵ Molar non functional cusps were found to have a steeper cuspal incline. As the cuspal inclines are the guiding planes for lateral excursive movements for group function occlusal relationships, these cusps may be subjected to greater occlusal forces. If other teeth in the arch have been restored with flatter cuspal inclines, then the steeper cusps are further exposed.⁴⁶ Over-carving of a restoration during placement, with loss of appropriate occlusal contact can result in the extrusion of a tooth, altering the cusp-fossae relationship and resulting in fracture of the non-functional cusp. However, the fracture of cusps, whether functional or non-functional, is primarily associated with large intra-coronal restorations and carious lesions.^{27,42,47}

CLINICAL SYMPTOMS

The clinical signs and symptoms may vary according to the




Fig. 4A. A split tooth where the mesio-distal fracture has resulted in separation of the segments. B. A radiograph of the tooth in Fig. 4A where the fracture is clearly seen. C. Separation of the fragments resulting in the split tooth is shown (arrow).




Fig. 5A. A deep and narrow periodontal defect is located with a probe on the buccal aspect of the right mandibular incisor. B. Surgical exposure confirms the presence and extent of the vertical root fracture (arrows).

position and extent of the incomplete fracture.^{13,43,48} Classically, the symptoms related to these teeth are pain on biting and sensitivity to thermal changes, particularly cold.^{2,3,11,13,18,31,49,50} Pain associated with the release of pressure, 'rebound pain' is also a consistent finding.^{12,13,18} Occasionally, there is sensitivity

to sweets.^{3,51} A chronic pulpitis with no clinical symptoms can exist as a result of microleakage of bacterial by-products and toxins. Pulpal and periodontal symptoms may occur when the fracture extends to involve the pulp.^{16,53}

DIAGNOSIS

A provisional diagnosis can generally be attained by a thorough history of the complaint. Early diagnosis is important, as restorative intervention can limit propagation of the fracture, subsequent microleakage and involvement of the pulpal or periodontal tissues, or catastrophic failure of the cusp.⁵⁴ The ease of diagnosis will vary according to the position and extent of the fracture.^{13,43} Dentin fractures are not generally evident radiographically, although radiographs are necessary to assess for caries, periapical status and the presence of periodontal lesions.^{13,31} Rubber dam isolation of the suspected tooth, and the application of cold or hot water are recommended. Once the tooth is identified, the offending cusp can be located by controlled wedging so as to load test individual cusps.^{2,13,31,43} A "Tooth Slooth"[®] is an appropriate instrument. When the tooth and cusp have been identified, the tooth can be anesthetized and all restorations removed to allow a thorough visual inspection so as to identify the position and extent of the fracture. The use of dyes, microscopes and transillumination are useful guides. Pulp sensibility testing of the tooth may be indicative of pulpal pathology. A tooth with an incomplete fracture may not be tender to percussion in a tooth with a healthy pulp.^{13,31}

MECHANISM OF PAIN

The character, duration and the stimuli of pain has important implications for both diagnosis and treatment.⁵⁵ An understanding of the mechanism of pain will often aid in assessment of the extent and direction of the crack. Luebke³² suggested the following terms to diagnose pain from a cracked tooth:

1. Dentin pain - A brief, sharp twinge.
2. Pulpal pain - The deep, demanding, radiating pain precipitated by thermal shock to an inflamed pulp. The pain at times may be spontaneous.
3. Periodontal pain - The aggravating throbbing of a sore tooth.

The pain associated with an incomplete fracture of a cusp is

generally accepted to be due to the rapid movement of dentin fluid in the dentin tubules according to the "Hydrodynamic theory of dentin sensitivity" as proposed and investigated by Brännström.^{16,18} Thermal changes, air, evaporation, osmotic stimuli such as sucrose, and increases in hydrostatic pressure caused by cuspal flexure as a result of occlusal forces can all act as stimuli for the rapid movement of dentin fluid. This movement stimulates A-delta nerve fibers in the vicinity of the odontoblastic processes and the pulp-dentin border, resulting in a sharp pain of short duration indicative of a vital tooth. Rebound pain, indicative of a vital tooth, is similarly explained when the pressure is released from the cusp as the tooth is free of the occlusion.^{16,18} When bacterial toxins have infiltrated the pulp, "hyperalgesia" can result. With this condition A-delta fibers are stimulated producing a sharp pain of short duration at what appears as a lower threshold than normal. The pain is due to the rapid movement of dentin fluid and probably a result of slight pulpal inflammation. During inflammation, the stimulation threshold of the A-delta fibers is lowered.⁵⁶

A second type of pulpal pain is produced by the stimulation of C-fibers as a response to inflammation, heat and mechanical deformation. A dull, poorly localized ache is often the result.⁵⁷ Alternatively, the pain can be a dull, aching pain with a continuous throbbing nature, or arise spontaneously and last for minutes or hours.⁵⁵ The C-fibers are activated by inflammatory mediators as a result of pulpal inflammation or prolonged application of heat.⁵⁸

A tooth with a painful pulpitis can present with a severe, sharp pain, indicative of A-delta fiber activation followed by a prolonged, dull ache that radiates throughout the jaw, indicative of C-fiber activation as well.⁵⁵ The C-fibers are resistant to tissue anoxia and can remain responsive long after the A-delta fibers.⁵⁹ A tooth with an incomplete fracture exhibiting C-fiber activation is strongly suggestive of pulpal damage and may require root canal treatment.

VERTICAL ROOT FRACTURE

The clinical presentation of a vertical root fracture is variable. Teeth with vertical root fractures often present with a history of discomfort and localized chronic inflammation. Patients may complain of a bad taste and pain on biting. If swelling is present it is generally broad-based and any sinus tract is located in or close to the attached gingiva rather than in the apical area. Double or multiple sinus tracts are common.⁷ A common feature of vertically root fractured teeth is the presence of a narrow periodontal pocket adjacent the fracture. Deep probing in two positions on opposite sides is almost pathognomonic for the presence of a fracture. The probing pattern for a tooth with a vertical root fracture is different from that seen with teeth with periodontal disease, where the pocketing is fairly consistent in depth around a large portion of the tooth.^{7,60}

The radiographic appearance of teeth with vertical root fractures is variable dependent on the angulation of X-ray beam in relation to the plane of the fracture and the degree of separation of the fragments. When separation of the root fragments has occurred, the root fracture is clearly visible. Alternatively, the radiographic image may show fracture lines along the root or root fillings, a space beside a root filling or

post, double images, radiolucent halos, unexplained bifurcation bone loss, J-shaped radiographic appearance, step-like bone defects, a widening of the periodontal ligament space, isolated horizontal bone loss in posterior teeth, V-shaped diffuse bone loss on roots of posterior teeth or dislodgement of a retrograde filling material.^{7,60}

While clinical and radiographic signs give a reasonably clear indication of the presence of a fracture, direct observation of the fracture is often required to confirm the presence of a fracture in many instances. This may involve a surgical approach and the use of transillumination is a useful diagnostic aid.

MANAGEMENT OF CRACKED TEETH

Coronal fracture

Management of cracked teeth should involve recognition of predisposing factors, recognition of signs and symptoms and the provision of adequate restorations that protect the tooth from fracture.⁴⁷ Early diagnosis is most important in the management of incomplete fracture so as to limit the propagation of the crack, subsequent microleakage and involvement of the pulpal and periodontal tissues.^{2,3,13,54} The treatment requirement of a cracked tooth is dependent on the position and extent of the fracture.^{13,43,61} An assessment of the stimuli, character and duration of the pain is also an influential guide for treatment.⁵⁵ As discussed, Luebke³² suggested pain from a cracked tooth be considered as dentin, pulpal or periodontal in character.

Cracks that enter the pulp indicate the need for root canal treatment though Bader *et al*⁶² reported that the majority of tooth fractures do not result in either pulp or tooth loss and can be managed successfully in a single visit using direct restorative materials. A multi-disciplinary approach involving endodontic, periodontic, orthodontic, prosthodontic and surgical intervention may be required.⁷ Fractures that involve the periodontal attachment may require extraction, though hemisection or root amputation may be appropriate for some multi-rooted teeth.^{7,63} However, teeth with cracks that are intraosseous with periodontal type pain often involving the mesial and distal aspects of the tooth and the cavity floor have a hopeless prognosis.^{64,65} A decision flow chart for the different classifications of cracked teeth can be seen in Fig 6.

Gutmann & Rakusin⁶⁵ suggested that treatments consist of an initial investigative and sedative stage followed by definitive treatment and restoration. Initial treatment involves the removal of all existing restorations to fully assess the extent of the fracture. Transillumination is a useful guide.³¹ In the initial diagnostic phase, the use of copper or stainless steel bands,^{13,43,49} stainless steel crowns,⁴⁸ and acrylic resin crowns⁶⁷ have been advocated. Placement of a sedative temporary restoration is not advised as this approach does not stabilize the fracture leaving the tooth susceptible for further extension of the crack.⁶⁸

In the absence of irreversible pulpitis, many techniques have been described to bind or remove the fracture so as to prevent flexure of the cusp, crack propagation and bacterial microleakage. Definitive treatment has included pin retained amalgams,^{61,64} bonded amalgams,^{52,69} bonded composites,⁷⁰⁻⁷³ cusp overlay restorations,⁷⁴⁻⁷⁶ and full coverage crowns.^{12,13,43,67}




Fig. 6. Treatment flow chart for AAE classification of cracked teeth. Adapted from Abbott.⁶⁶

Teeth restored with cuspal amalgam overlays had fracture energies, measured as the force required to fracture, equal to that of an intact tooth whereas gold crowns increased the fracture energy by more than three-fold.⁷⁷

Clark & Caughman⁶⁴ have categorized the prognosis of cracked teeth as excellent, good, poor and hopeless.

1. Excellent: (a) Cuspal fracture confined within the dentin that angles from the facio-pulpal or linguo-pulpal line angle of a cusp to the cemento-enamel junction or slightly below. (b) Horizontal fracture of a cusp not involving the pulp.
2. Good: A coronal vertical fracture that runs mesio-distally into the dentin but not into the pulp.
3. Poor: A coronal vertical fracture that runs mesio-distally into the dentin and pulp but is confined to the crown.
4. Hopeless: A coronal vertical fracture that runs mesio-distally through the pulp and extends into the root.

The provision of an acrylic splint is recommended for prevention of further fractures in patients with parafunctional occlusal activity or a history of incomplete fracture in other teeth.⁵⁰

Vertical root fracture

Single-rooted teeth that are fractured should be extracted as soon as is practical to prevent further bone loss. Multi-rooted teeth can often be successfully treated by resecting the fractured root, either by root amputation or hemisection.⁷⁸ Studies of root resected teeth have reported 5-year retention rates of 94%⁷⁹ and 10-year retention rates of 68%.⁸⁰ However, the desire to retain part of a root fractured tooth should be carefully considered against extraction and replacement with a denture, bridge or implant.

SUMMARY

The clinical features, diagnosis and management of cracked teeth have been reviewed. Fractures may initiate from coronal tooth structure or from within the root. There are many terminologies and classifications in the literature for cracked teeth which can be as confusing as the array of clinical symptoms that are associated with this condition. The term “cracked tooth syndrome” is misleading as there are a range of symptoms that do not form a distinct and reliable pattern. Symptoms will vary with teeth that have healthy pulps, for

teeth with inflamed or necrotic pulps, and for teeth that have been root filled.

The American Association of Endodontists has classified five specific variations of cracked teeth; craze line, fractured cusp, cracked tooth, split tooth, and vertical root fracture. Furthermore, the character, duration and the stimuli of pain have important implications for both diagnosis and treatment. Diagnostically, it is important to differentiate the differences between dentin, pulpal and periodontal pain before treatment is commenced. Early diagnosis is most important in the treatment of cracked teeth to limit the propagation of the crack. A decision flow chart indicating the treatment options available has been presented to clarify the cracked tooth conundrum.

a. Professional Results, Inc., Laguna Niguel, CA, USA.

Acknowledgements: To Professor Geoffrey Heithersay and Drs. Tom Berekally and John Abbott for their helpful comments on this manuscript, and to Associate Professor Alex Moule and Drs. Tom Berekally and Fab Damiani for kindly providing clinical photographs.

Dr. Kahler is a PhD candidate at the University of Sydney, Australia, a Senior Lecturer at the University of Queensland and Endodontist in Brisbane, Australia.

References

- Bender IB, Freedland JB. Adult root fracture. *J Am Dent Assoc* 1983;107:413-419.
- Cameron CE. The cracked tooth syndrome. *J Am Dent Assoc* 1964;68:405-411.
- Hiatt WH. Incomplete crown-root fracture in pulpal-periodontal disease. *J Periodontol* 1973;44:369-379.
- Walton RE, Michelich RJ, Smith G.N. The histopathogenesis of vertical root fractures. *J Endod* 1984;10:48-56.
- Goel VK, Khera SC, Gurusami S, Chen RCS. Effect of cavity depth on stresses in a restored tooth. *J Prosthet Dent* 1992;67:174-183.
- Arola D, Huang MP, Sultan MB. The failure of amalgam dental restorations due to cyclic fatigue crack growth. *J Mater Sci Mater Med* 1999;10:319-327.
- Pitts DL, Natkin E. Diagnosis and treatment of vertical root fractures. *J Endod* 1983;9:338-346.
- Meister F, Lommel TJ, Gerstein H. Diagnosis and possible causes of vertical root fractures. *Oral Surg* 1980;49:243-253.
- Holcomb JQ, Pitts DL, Nicholls JI. Further investigation of spreader loads required to cause vertical root fracture during lateral condensation. *J Endod* 1987;13:277-284.
- Murgel CAF, Walton RE. Vertical root fracture and dentine deformation in curved roots: The influence of spreader design. *Endod Dent Traumatol* 1990;6:273-278.
- Gibbs JW. Cuspal fracture odontalgia. *Dent Digest* 1954;60:158-160.
- Cameron CE. The cracked tooth syndrome: Additional findings. *J Am Dent Assoc* 1976;93:971-975.
- Ehrmann EH, Tyass MJ. Cracked-tooth syndrome: Diagnosis, treatment and correlation between symptoms and post-extraction findings. *Aust Dent J* 1990;35:105-102.
- Geurtsen W, Schwarze T, Günay H. Diagnosis, therapy, and prevention of the cracked tooth syndrome. *Quintessence Int* 2003;34:409-417.
- Ellis SGS. Incomplete tooth fracture-proposal for a new definition. *Br Dent J* 2001;190:424-428.
- Brännström M. *Dentin and pulp in restorative dentistry*. London: Wolfe Medical Publications Ltd., 1982;47-63.
- Kahler B, Stenzell D, Moule A. Bacterial contamination of cracks in symptomatic vital teeth. *Aust Endod J* 2000;26:115-117.
- Brännström M. The hydrodynamic theory of dentinal pain: Sensation in preparations, caries, and the dentinal crack syndrome. *J Endod* 1986;12:453-457.
- Thoma KH. *Oral pathology*. 4th ed. St. Louis: Mosby, 1954.
- Ritchie B, Mendenhall R, Orban B. Pulpitis resulting from incomplete tooth structure. *Oral Surg Oral Med Oral Pathol* 1957;10:665-670.
- Down CH. The treatment of permanent incisor teeth of children following traumatic injury. *Aust Dent J* 1957;2:9.
- Sutton PRN. Transverse crack lines in permanent incisors of Polynesians. *Aust Dent J* 1961;6:144-150.
- Sutton PRN. Greenstick fracture of the tooth crown. *Br Dent J* 1962;112:362-363.
- Wiebusch FB. Hairline fracture of a cusp: Report of a case. *J Can Dent Assoc* 1972;38:192-194.
- Talim ST, Gohil KS. Management of coronal fractures of permanent posterior teeth. *J Prosthet Dent* 1974;31:172-178.
- Silvestri AR. The undiagnosed split-root syndrome. *J Am Dent Assoc* 1976;92:930-935.
- Maxwell EH, Braly BV. Incomplete tooth fracture: Prediction and prevention. *CDA J* 1977;5:51-55.
- Andreasen JO. *Traumatic injuries of the teeth*. 3rd ed. Copenhagen: Munksgaard, 1994.
- Caufield JB. Hairline tooth fracture: A clinical case report. *J Am Dent Assoc* 1981;102:501-502.
- Johnson R. Descriptive classification of traumatic injuries to the teeth and supporting structures. *J Am Dent Assoc* 1981;102:195-197.
- Abou-rass M. Crack lines: The precursors of tooth fracture. Their diagnosis and treatment. *Quintessence Int* 1983;14:437-444.
- Luebke RG. Vertical crown-root fractures in posterior teeth. *Dent Clin North Am* 1984;28:883-894.
- Kruger BF. Cracked cusp syndrome. *Aust Dent J* 1984;29:55.
- Williams J. Incomplete vertical tooth fracture. *J Mass Dent Soc* 1988;37:13-20.
- Löst C, Bengel W, Hehner B. Tooth infraction. Incomplete tooth fracture. A review of various aspects of the disease with case reports. *Schweiz Monatsschr Zahnmed* 1989;99:1033-1040.
- Schweitzer JL, Gutmann JL, Bliss RQ. Odontiatrogenic tooth fracture. *Int Endod J* 1989;22:64-74.
- Zuckerman GR. The cracked tooth. *NY State Dent J* 1998;126:30-35.
- Pruden WH. Treatment of the cracked tooth. *J N J Dent Assoc* 1971;42:22-23.
- Clark DJ, Sheets CG, Paquette JM. Definitive diagnosis of early enamel and dentin cracks based on microscopic evaluation. *J Esthet Restor Dent* 2003;15:391-401.
- <http://www.aae.org/dentalpro/colleagueneews.htm>
- Snyder DE. The cracked-tooth syndrome and fractured posterior cusp. *Oral Surg* 1976;41:698-704.
- Eakle WS. Increased fracture resistance of teeth: Comparison of five bonded composite resin systems. *Quintessence Int* 1986;17:17-20.
- Geurtsen W. The cracked-tooth syndrome: Clinical features and case reports. *Int J Periodontol Rest Dent* 1992;12:395-405.
- Lagouvardos P, Sourai P, Douvitisas C. Coronal fractures in posterior teeth. *Oper Dent* 1989;14:28-32.
- Cavel WT, Kelsey WP, Blankenau RJ. An *in vivo* study of cuspal fracture. *J Prosthet Dent* 1985;53:38-41.
- Khera SC, Carpenter CW, Vetter JD, Staley RN. Anatomy of cusps of posterior teeth and their fracture potential. *J Prosthet Dent* 1990;64:139-147.
- Braly BV, Maxwell EH. Potential for tooth fracture in restorative dentistry. *J Prosthet Dent* 1981;45:411-414.
- Chong BS. Bilateral cracked teeth: A case report. *Int Endod J* 1989;22:193-196.
- Homewood CI. Cracked tooth syndrome. Incidence, clinical findings and treatment. *Aust Dent J* 1998;43:217-222.
- Zimet PO. Cracked tooth syndrome. *Aust Endod J* 1998;24:33-37.
- Stanley HR. The cracked tooth syndrome. *J Am Acad Gold Foil Oper* 1968;11:36-47.
- Trushkowsky R. Restoration of a cracked tooth with a bonded amalgam. *Quintessence Int* 1991;22:397-400.
- Bergenholtz G. Pathogenic mechanisms in pulpal disease. *J Endod* 1990;16:98-101.
- Agar JR, Weller RN. Occlusal adjustment for initial treatment and prevention of cracked-tooth syndrome. *J Prosthet Dent* 1988;60:145-147.
- Figdor D. Pain of dentinal and pulpal origin. A review for the clinician. *Ann R Coll Dent Surg* 1994;12:131-142.
- Trowbridge HO. Review of dental pain-histology and physiology. *J Endod* 1986;12:445-452.
- Jyväsjarvi E, Kniffki KD. Afferent C fibre innervation of cat tooth pulp: Confirmation by electrophysiological methods. *J Physiol* 1989;411:663-675.
- Jyväsjarvi E, Kniffki KD, Mengel MKC. Functional characteristics of afferent C fibres from tooth pulp and periodontal ligament. In: Hammann W, Iggo A. *Progress in brain research*. London: Elsevier, 1988;74:237-245.
- Närhi MVO. The characteristics of intradental sensory units and their

- responses to stimulation. *J Dent Res* 1985;64 (Sp Is):564-571.
60. Moule AJ, Kahler B. Diagnosis and management of teeth with vertical root fractures. *Aust Dent J* 1999;44:75-87.
 61. Silvestri AR, Singh I. Treatment rationale of fractured posterior teeth. *J Am Dent Assoc* 1978;97:806-810.
 62. Bader JD, Shugars DA, Sturdevant JR. Consequences of posterior cusp fracture. *Gen Dent* 2004;63:128-131.
 63. Burke FJT. Tooth fracture *in vivo* and *in vitro*. *J Dent* 1992;20:131-139.
 64. Clark LL, Caughman WF. Restorative treatment for the cracked tooth. *Oper Dent* 1984;9:136-142.
 65. Gutmann JL, Rakusin H. Endodontic and restorative management of incompletely fractured molar teeth. *Int Endod J* 1994;27:343-348.
 66. Abbott, P. Endodontics and dental traumatology. An overview of modern endodontics. 1999
 67. Guthrie C, Difiore PM. Treating the cracked tooth with a full crown. *J Am Dent Assoc* 1991;122:71-73.
 68. Ailor JE. Managing incomplete fractures. *J Am Dent Assoc* 2000;131:1168-1174.
 69. Bearn DR, Saunders EM, Saunders WP. The bonded amalgam restoration. A review of the literature and report of its use in the treatment of four cases of cracked-tooth syndrome. *Quintessence Int* 1994;25:321-326.
 70. Eakle WS. Reinforcement of fractured posterior teeth with bonded composite restorations. *Quintessence Int* 1985;16:481-482.
 71. Baxter PW. Management of vertical fractures of posterior teeth with composite resin. *Br Dent J* 1987;162:219-220.
 72. Hansen EK. *In vivo* cusp fracture of endodontically treated premolars restored with MOD amalgam or MOD resin fillings. *Dent Mater* 1988;4:169-173.
 73. Burke FJT, Wilson NHF, Watts DC. Fracture resistance of teeth restored with indirect composite restorations: The effect of alternative luting procedures. *Quintessence Int* 1994;25:269-275.
 74. Hansen EK, Asmussen E, Christiansen NC. *In vivo* fractures of endodontically treated posterior teeth restored with amalgam. *Endod Dent Traumatol* 1990;6:49-55.
 75. Linn J, Messer HH. Effect of restorative procedures on the strength of endodontically treated molars. *J Endod* 1994;20:479-485.
 76. Panitvisai P, Messer HH. Cuspal deflection in molars in relation to endodontic and restorative procedures. *J Endod* 1995;21:57-61.
 77. Salis SG, Hood JAA, Kirk EEJ, Stokes ANS. Impact-fracture energy of human premolar teeth. *J Prosthet Dent* 1987;58:43-48.
 78. Korte PF, Carr JG, Cohen J. Vertical root fracture and its relationship to the periodontium. *J Mich Dent Assoc* 1980;62:387-389.
 79. Langer B, Stein S, Wagenberg B. An evaluation of root resections. A ten year study. *J Periodontol* 1981;52:719-722.
 80. Buhler H. Evaluation of root-resected teeth. Results after 10 years. *J Periodontol* 1988;59:805-810.